

TABLE 7. ATP requirement for biosyntheses during bacterial growth

Carbon source	ATP requirement (mol of ATP/g of cells)	$Y_{ATP}^m$ (g of cells/mol of ATP)	Reference
Glucose	0.037	27	626
Lactate <sup>-</sup>	0.075	13.4	626
Malate <sup>2-</sup>	0.065	15.4	626
Acetate <sup>-</sup> + CO <sub>2</sub>	0.065	15.4	129, 642
Acetate <sup>-</sup>	0.1	10	626
CO <sub>2</sub>	0.2	5	177

129. Decker, K., K. Jungermann, and R. K. Thauer.  
 1970. Energy production in anaerobic organism. *Angew. Chem. Int. Ed. Engl.* 9:138-158.
177. Forrest, W. W., and D. J. Walker. 1971. The generation and utilization of energy during growth. *Adv. Microbiol. Physiol.* 5:213-274.
626. Stouthamer, A. H. 1973. A theoretical study on the amount of ATP required for synthesis of microbial cell material. *Antonie van Leeuwenhoek J. Microbiol. Serol.* 39:545-565.
642. Thauer, R. K. 1967. Der Energiestoffwechsel von *Clostridium kluyveri*. Ph.D. thesis, The University of Freiburg, Germany.