

Table 1. Frequency of Ig-producing cells in EBV-infected BM^a

BM sample	Frequency $\times 10^{-4}$	IgM Statistics (p)	Isotype distribution (%)	Frequency $\times 10^{-4}$	IgG Statistics (p)	Isotype distribution (%)	Frequency $\times 10^{-4}$	IgA Statistics (p)	Isotype distribution (%)
1	1.2 ± 0.4	0.39	(9.7)	7.4 ± 1.1	0.31	(59.7)	3.8 ± 0.7	0.79	(30.6)
2	1.8 ± 0.6	0.55	(10.8)	8.1 ± 2.1	0.63	(48.5)	6.8 ± 1.9	0.71	(40.7)
3	2.8 ± 0.3	0.99	(15.6)	10.1 ± 1.2	0.87	(56.4)	5.0 ± 7.3	0.44	(27.9)
4	3.4 ± 0.4	0.55	(20.8)	7.6 ± 1.4	0.87	(46.6)	5.3 ± 1.3	0.97	(32.5)
5	7.7 ± 0.9	0.55	(31.4)	8.7 ± 0.9	0.78	(35.5)	8.1 ± 0.8	0.50	(33.1)
6	6.8 ± 0.8	0.64	(36.4)	6.5 ± 0.8	0.92	(34.7)	5.4 ± 0.7	0.87	(28.9)
7	1.2 ± 0.2	0.17	(20.0)	3.1 ± 0.5	0.98	(51.7)	1.7 ± 0.3	0.95	(28.3)
8	3.1 ± 0.3	0.93	(48.4)	1.6 ± 0.2	0.45	(25.0)	1.7 ± 0.2	0.98	(26.6)
9	2.3 ± 0.4	0.18	(17.3)	5.0 ± 0.6	0.23	(37.6)	6.0 ± 0.8	0.78	(26.3)
10	8.2 ± 1.7	0.32	(46.8)	4.7 ± 1.0	0.14	(26.8)	4.6 ± 0.10	0.14	(26.3)
Average	3.9 ± 2.5	0.53	(26.0)	6.3 ± 2.5	0.62	(42.0)	4.8 ± 1.93	0.17	(32.0)

a) Frequency of cells producing a given Ig isotype as calculated by the maximum likelihood procedure.