

Table S4. Half-life of pre-mRNA in mouse embryonic fibroblasts (3T3)

Gene	Location	Fibroblasts		
		Basal	TNF (30 min)	TNF (6 h)
Housekeeping				
<i>Actb</i>	In 2	4.6 ± 0.2	3.3 ± 0.4	4.4 ± 1.2
Group I				
<i>Cxcl1</i>	In 1	2.7 ± 0.3	2.0 ± 0.1	2.1 ± 0.3
<i>Nfkbia</i>	In 5	3.3 ± 0.7	2.9 ± 0.3	3.2 ± 0.2
<i>Tnfaip3</i>	In 2	5.6 ± 1.5	4.2 ± 0.3	3.2 ± 0.7
<i>Fos</i>	In 1	—	2.9 ± 0.8	4.0 ± 3.7
Group II				
<i>Cxcl10</i>	In 1	—	5.1 ± 1.5	4.3 ± 1.1
<i>Icam1</i>	In 2	—	7.4 ± 6.2	6 ± 4
<i>Nfkbie</i>	In 5	9.0 ± 4.2	13	15
Group III				
<i>Saa3</i>	In 3 (3')	6.8 ± 1.3	4.2 ± 0.4	3.7 ± 0.2
<i>Mmp3</i>	In 8	—	10 ± 2	8.3 ± 3.2
<i>Ccl5</i>	In 2	12 ± 5	20 ± 12	10.5 ± 3.9
<i>Gbp2</i>	In 1	—	>30	16 ± 3
<i>Ifit1</i>	In 1	—	>30	24 ± 11

For methods and notes, see Table S3.