

Table 1. Reported or derived data on the lymphocyte content of different lymphoid organs and tissues in normal young adult rats of 200 g weight

Authors		Kindred (1942) × 10 ⁹	Monden (1958) × 10 ⁹	Everett (1964) × 10 ⁹	Ford (1969) × 10 ⁹	Average × 10 ⁹
Lymph nodes	(absolute number) (/kg body weight)	2.1 10.5	1.7 8.5			1.9 9.5
Spleen	(abs.) (/kg)	0.2 1.0	1.1 5.5	0.3 1.5	0.7 2.8	0.6 3.0
Peyer's patches	(abs.) (/kg)	0.03 0.15	0.4 2.0			0.22 1.1
Thymus	(abs.) (/kg)	1.0 5.0	1.1 5.5	1.8 9.0		1.3 6.5
Total lymphocytes in lymphoid tissue	(abs.) (/kg)	3.33 16.60	4.3 21.5			4.0 20.0
Bone marrow	(abs.) (/kg)	0.9 4.5		0.7 3.5		0.8 4.0
Blood	(abs.) (/kg)	0.08 0.4	0.13 0.7			0.11 0.55
Total lymphocytes in lymphoid tissue, bone marrow and blood	(abs.) (/kg)					4.9 24.5

Everett, N. B., Caffrey, R. W., Rieke, W. O.: Recirculation of lymphocytes. Ann. N. Y. Acad. Sci. 118, 887-897 (1964)
 Ford, W. L.: The kinetics of lymphocyte recirculation within the rat spleen. Cell Tiss. Kinet. 2, 171-191 (1969)

Kindred, J. E.: A quantitative study of the hemopoietic organs of young adult albino rats. Amer. J. Anat. 71, 207-243 (1942)

Monden, Y.: Quantitative evaluation of total cellular number and cellular density in the thymolymphatic organs of young adult albino rats by means of DNA determination. Okajimas Folia anat. Jap. 32, 193-206 (1959)