

15. **Harmsen, H., G. Raangs, T. He, J. Degener, and G. Welling.** 2002. Extensive set of 16S rRNA-based probes for detection of bacteria in human feces. *Appl. Environ. Microbiol.* **68**:2982–2990.
18. **Hold, G., S. Pryde, V. Russell, E. Furrrie, and H. Flint.** 2002. Assessment of microbial diversity in human colonic samples by 16S rDNA sequence analysis. *FEMS Microbiol. Ecol.* **39**:33–39.
22. **Matsuki, T., K. Watanabe, J. Fujimoto, Y. Miyamoto, T. Takada, K. Matsumoto, H. Oyaizu, and R. Tanaka.** 2002. Development of 16S rRNA-gene-targeted group-specific primers for the detection and identification of predominant bacteria in human feces. *Appl. Environ. Microbiol.* **68**:5445–5451.
25. **Nagashima, K., T. Hisada, S. Maremi, and J. Mochizuki.** 2003. Application of new primer-enzyme combinations to terminal restriction fragment length polymorphism profiling of bacterial populations in human feces. *Appl. Environ. Microbiol.* **69**:1251–1262.
32. **Suau, A., R. Bonnet, M. Sutren, J.-J. Godon, G. R. Gibson, M. D. Collins, and J. Dore.** 1999. Direct analysis of genes encoding 16S rRNA from complex communities reveals many novel molecular species within the human gut. *Appl. Environ. Microbiol.* **65**:4799–4807.
35. **Wang, R., S.-J. Kim, L. Robertson, and C. Cerniglia.** 2002. Development of a membrane-array method for the detection of human intestinal bacterial in fecal samples. *Mol. Cell. Probes* **16**:341–350.