

TABLE 3. *Estimates of contributions of VFA produced in different segments of digestive tract of various species to energy requirements of whole body*

| Species | Organ | % of Energy Requirements | Reference |
|-----------|------------------|--------------------------|-----------|
| Sheep | Rumen* | 70 | 51, 128 |
| | Cecum* | 5 | 110 |
| | Total hindgut† | 8 | 143, 294 |
| Cattle | Rumen* | 63 | 269 |
| | Total hindgut† | 9 | 269 |
| Rat | Cecum* | 5 | 330 |
| Porcupine | Cecum‡ | 16 | 164 |
| Beaver | Large intestine‡ | 19 | 144 |
| Rabbit | Cecum‡ | 12 | 145 |
| | Total hindgut* | 30 | 203, 221 |
| | Cecum* | 30 | 123 |
| Pony | Large intestine‡ | 11 | 160, 175 |
| | Total hindgut§ | 25 | 244 |
| Human | Large intestine† | 6-10 | 209 |

* In vivo isotope dilution. † Calculations based on amount of fiber and carbohydrate leaving ileum. ‡ In vitro incubation of gut contents. § Absorption into portal blood.

51. BERGMAN, E. N., R. S. REID, M. G. MURRAY, J. M. BROCKWAY, AND F. G. WHITELAW. Interconversions and production of volatile fatty acids in the sheep rumen. *Biochem. J.* 97: 53-58, 1965.
110. FAICHNEY, G. J. Production of volatile fatty acids in the sheep caecum. *Aust. J. Agric. Res.* 20: 491-498, 1969.
123. GLINSKY, M. J., R. M. SMITH, H. R. SPIRES, AND C. L. DAVIES. Measurement of volatile fatty acid production rates in the cecum of the pony. *J. Anim. Sci.* 42: 1465-1470, 1976.
128. GRAY, F. V., R. A. WELLER, A. F. PILGRIM, AND G. B. JONES. Rates of production of volatile fatty acids in the rumen. *Aust. J. Agric. Res.* 18: 625-634, 1967.

143. HOOVER, W. H. Digestion and absorption in the hindgut of ruminants. *J. Anim. Sci.* 46: 1789-1799, 1978.
144. HOOVER, W. H., AND S. D. CLARKE. Fiber digestion in the beaver. *J. Nutr.* 102: 9-15, 1972.
145. HOOVER, W. H., AND R. N. HEITMANN. Effects of dietary fiber levels on weight gain, cecal volume, and volatile fatty acid production in rabbits. *J. Nutr.* 102: 375-380, 1972.
160. IMOTO, S., AND S. NAMIOKA. VFA production in the pig large intestine. *J. Anim. Sci.* 47: 467-478, 1978.
164. JOHNSON, J. L., AND R. H. MCBEE. The porcupine cecal fermentation. *J. Nutr.* 91: 540-546, 1967.
175. KIM, K. I., N. G. BENEVENGA, AND W. H. GRUMMER. Lactase activity and VFA production in the caecum and colon of pigs fed a corn-soy or 40% whey diet. *J. Anim. Sci.* 46: 1648-1657, 1978.
203. MARTY, J., AND M. VERNAY. Absorption and metabolism of the volatile fatty acids in the hindgut of the rabbit. *Br. J. Nutr.* 51: 265-277, 1984.
209. MCNEIL, N. I. The contribution of the large intestine to energy supplies in man. *Am. J. Clin. Nutr.* 39: 338-342, 1984.
221. PARKER, D. S. The measurement of the production rates of volatile fatty acids in the caecum of the conscious rabbit. *Br. J. Nutr.* 36: 61-70, 1976.
244. RERAT, A., M. FISZLEWICZ, A. GIUSI, AND P. VAUGELADE. Influence of meal frequency on postprandial variations in the production and absorption of volatile fatty acids in the digestive tract of conscious pigs. *J. Anim. Sci.* 64: 448-456, 1987.
269. SICILIANO-JONES, J., AND M. R. MURPHY. Production of volatile fatty acids in the rumen and cecum-colon of steers as affected by forage: concentrate and forage physical form. *J. Dairy Sci.* 72: 485-492, 1989.
294. ULYATT, M. J., D. W. DELLOW, C. S. W. REID, AND T. BAUCHOP. Structure and function of the large intestine. In: *Digestion and Metabolism in the Ruminant*, edited by I. W. McDonald and A. C. I. Warner. Armidale, Australia: Univ. of New England, 1975, p. 119-133.
330. YANG, M. G., K. MANOHARAN, AND O. MICHELSEN. Nutritional contribution of volatile fatty acids from the cecum of rats. *J. Nutr.* 100: 545-550, 1970.