

TABLE 1. In Vitro Determination of the Effect of Coconut Water (CNW) or Physiological Saline (PS) on the TEG Parameters in 8 Healthy Volunteers

Parameter	Baseline	Units Citrated Plasma:Units CNW			Units Citrated Plasma:Units PS		
		10:2	10:4	10:10	10:2	10:4	10:10
SP (min)	9.2 ± 0.69	7.8 ± 0.80	7.9 ± 0.59	8.5 ± 0.59	9.3 ± 0.65	8.8 ± 0.55	9.1 ± 0.48
R (min)	9.9 ± 0.81	9.1 ± 0.61	8.3 ± 0.60	9.2 ± 0.63	10.2 ± 0.79	9.5 ± 0.53	9.9 ± 0.42
K (min)	3.5 ± 0.20	3.4 ± 0.23	4.2 ± 0.30	6.8 ± 0.53*	3.6 ± 0.25	4.0 ± 0.28	5.8 ± 0.24*
MA (mm)	54.7 ± 1.76	50.5 ± 1.77*	46.9 ± 2.00*	33.4 ± 1.46*	51.2 ± 1.20*	46.4 ± 1.21*	37.3 ± 1.22*
α (deg)	68.0 ± 1.34	65.9 ± 1.42	65.9 ± 1.14	56.4 ± 1.76*	66.5 ± 1.52	66.7 ± 1.42	59.0 ± 0.92*

NOTE. Data are expressed as mean ± SEM; paired t-test was performed to compare CNW with baseline or PS.

* $P < .05$ v baseline; there was no significant difference between CNW and PS.