

Table 2Percentage of constituents in rhizome with roots of *Rhodiola rosea* [34,51–53,74,75].

Constituents	Important markers	Percentage (%)
Phenylpropanoids	Rosarin	0.025
	Rosavin	0.208
	Rosin	0.121
Phenylethanol derivatives	Tyrosol	0.077
	Salidroside	0.155
Flavonoids	Catechins	0.165
Phenolic acids	Gallic acid, Caffeic acid, Chlorogenic acid	0.067

- [34] Z. Liu, Y. Liu, C. Liu, Z. Song, Q. Li, Q. Zha, C. Lu, C. Wang, Z. Ning, Y. Zhang, C. Tian, Lu A. The chemotaxonomic classification of *Rhodiola* plants and its correlation with morphological characteristics and genetic taxonomy, *Chem. Cent. J.* 7 (1) (2013) 118.
- [51] M. Du, J. Xie, Flavonol glycosides from *Rhodiola crenulata*, *Phytochemistry* 38 (February (3)) (1995) 809–810.
- [52] X. Li, S. Nakamura, H. Matsuda, M. Yoshikawa, Bioactive constituents from chinese natural medicines. XXIX. Monoterpene and monoterpene glycosides from the roots of *Rhodiola sachalinensis*, *Chem. Pharm. Bull. (Tokyo)* 56 (April (4)) (2008) 612–615.
- [53] A.B. Tayade, P. Dhar, M. Sharma, R.S. Chauhan, O.P. Chaurasia, R.B. Srivastava, Antioxidant capacities, phenolic contents, and GC/MS analysis of *Rhodiola imbricata* Edgew. root extracts from Trans-Himalaya, *J. Food Sci.* 78 (March (3)) (2013) C402–C410.
- [74] A. Adamczak, W. Buchwald, A. Gryszczyńska, Biometric features and content of phenolic compounds of roseroot (*Rhodiola rosea* L.), *Acta Soc. Bot. Pol.* 85 (3) (2016) 3500.
- [75] B. Kolodziej, D. Sugier, Influence of plants age on the chemical composition of roseroot (*Rhodiola rosea* L.), *Acta Sci. Pol. Hortorum Cultus* 12 (January (3)) (2013) 147–160.