

Table 3. Female breast and reproductive tract tumours

Reference	No.	Median LI % range	Median Ts (h) range	Median Tpot (days) range	Comment
Breast					
Rew (1991) ^{42,43}	75	4.2 (0.6–15.4)	8.7 (2.7–22.2)	8.2 (1.8–47.5)	BrdUrd 250 mg
Stanton (1996) ^{45,46}	84	3.2	12.0	12.5	BrdUrd 200 mg
Gynaecology					
Ovarian					
Erba (1994) ⁴⁷	55	6.1–7.2	14.7	12.5	BrdUrd 250 mg
Cervix					
Bolger (1996) ^{48,49}	120	9.8 (6.7–14.5)	12.8 (11.1–14.9)	4.0 (3.1–6.3)	BrdUrd 200 mg
(Squamous)		Interquartile	Interquartile	Interquartile	
Tsang (1995) ⁵⁰	39			5.5	BrdUrd 200 mg
(Adenocarc)					
Tsang (1995) ⁵⁰	7			6.6	BrdUrd 200 mg
Ovarian metastases					
Rew (1991) ²²	2	5.1, 5.4	12.5, 13.9	3.5, 6.5	BrdUrd 250 mg

LI: labelling index; Ts: S-phase duration; Tpot: potential doubling time.