

Table S1: pH and electrical potential ($\Delta\phi$) in each of the eight cellular compartments included in Recon 1. *Variability in peroxisomal pH and $\Delta\phi$ was accounted for by adding uncertainty to standard transformed Gibbs energy estimates for peroxisomal metabolites and reactions (see Sections A.3.1 and A.3.2).

Compartment	pH	$\Delta\phi$ (mV)	References
Cytosol and nucleus	7.20	0	(4)
Extracellular fluid	7.40	30	(7), (21, 22)
Golgi apparatus	6.35	0	(6), (25, 26)
Lysosomes	5.50	19	(6), (28)
Mitochondria	8.00	-155	(67), (20)
Endoplasmic reticulum	7.20	0	(6), (26)
Peroxisomes*	7.00 ± 1.2	12 ± 74	(11, 12, 13, 14, 15), (16, 18)

4. Buckler, K. J., and R. D. Vaughan-Jones, 1990. Application of a new pH-sensitive fluoroprobe (carboxy-SNARF-1) for intracellular pH measurement in small, isolated cells. *Pflügers Arch.* 417:234–239.
6. Paroutis, P., N. Touret, and S. Grinstein, 2004. The pH of the secretory pathway: measurement, determinants, and regulation. *Physiology* 19:207–215.
7. Street, D., J. Bangsbo, and C. Juel, 2001. Interstitial pH in human skeletal muscle during and after dynamic graded exercise. *J. Physiol.* 537:993–998.
11. Dansen, T. B., K. W. A. Wirtz, R. J. A. Wanders, and E. H. W. Pap, 1999. Peroxisomes in human fibroblasts have a basic pH. *Nat. Cell. Biol.* 2:51–53.
12. Douma, A. C., M. Veenhuis, G. J. Sulter, and W. Harder, 1987. A proton-translocating adenosine triphosphatase is associated with the peroxisomal membrane of yeasts. *Arch. Microbiol.* 147:42–47.
13. Jankowski, A., J. H. Kim, R. F. Collins, R. Daneman, P. Walton, and S. Grinstein, 2001. In situ measurements of the pH of mammalian peroxisomes using the fluorescent protein pHluorin. *J. Biol. Chem.* 276:48748–48753.
14. Nicolay, K., M. Veenhuis, A. C. Douma, and W. Harder, 1987. A 31 P NMR study of the internal pH of yeast peroxisomes. *Arch. Microbiol.* 147:37–41.
15. van Roermund, C. W. T., M. de Jong, L. IJlst, J. van Marle, T. B. Dansen, R. J. A. Wanders, and H. R. Waterham, 2004. The peroxisomal lumen in *Saccharomyces cerevisiae* is alkaline. *J. Cell. Sci.* 117:4231–4237.
16. Antonenkov, V. D., and J. K. Hiltunen, 2006. Peroxisomal membrane permeability and solute transfer. *Biochim. Biophys. Acta, Mol. Cell. Res.* 1763:1697–1706.
18. Antonenkov, V. D., and J. K. Hiltunen, 2011. Transfer of metabolites across the peroxisomal membrane. *Biochim. Biophys. Acta*.
20. Cortese, J. D., 1999. Rat liver GTP-binding proteins mediate changes in mitochondrial membrane potential and organelle fusion. *Am. J. Physiol.* 276:C611–C620.
21. Hoek, J. B., D. G. Nicholls, and J. R. Williamson, 1980. Determination of the mitochondrial protonmotive force in isolated hepatocytes. *J. Biol. Chem.* 255:1458–1464.
22. Nobes, C. D., G. C. Brown, P. N. Olive, and M. D. Brand, 1990. Non-ohmic proton conductance of the mitochondrial inner membrane in hepatocytes. *J. Biol. Chem.* 265:12903–12909.
25. Schapiro, F. B., and S. Grinstein, 2000. Determinants of the pH of the Golgi complex. *J. Biol. Chem.* 275:21025–21032.
26. Wu, M. M., J. Llopis, S. Adams, J. M. McCaffery, M. S. Kulomaa, T. E. Machen, H. P. Moore, and R. Y. Tsien, 2000. Organelle pH studies using targeted avidin and fluorescein-biotin. *Chem. Biol.* 7:197–209.
28. Koivusalo, M., B. E. Steinberg, D. Mason, and S. Grinstein, 2011. In situ measurement of the electrical potential across the lysosomal membrane using FRET. *Traffic* 12:972–982.

67. Wilson, L. G., and R. S. Bandurski, 1958. Enzymatic reactions involving sulfate, sulfite, selenate, and molybdate. *J. Biol. Chem.* 233:975–981.